

TUM-HEP-493/02
hep-ph/0211407

CP-ODD PHASES @ NLC: A PROJECT OUTLINE

B.GAISSMAIER¹

Physik Dept., TU München, James Franck Str., D-85748 Garching, Germany

Abstract

Low-energy results from measurements of leptonic dipole operators are used to derive constraints on phases of the MSSM. After rediscovering older bounds on these phases, we try to investigate the impact of these possibly non-vanishing CP-odd phases on the measurement of CP-even cross-sections at the next leptonic collider.

1 Introduction

The most general MSSM provides a rather big number of complex phases, most of which originate from the soft breaking terms. Of course, the possible ranges of these phases are restricted by the current measurements of leptonic dipole operators such as d_e [1] or a_μ [2],[3]. Fortunately, theory can provide at least one scenario (cancellation), in which the bounds for the dipole operators are respected without too small values for the phases [4],[5]. Once assumptions about the mechanism of supersymmetry breaking are made, most soft breaking parameters can be computed from their input parameter using the RGE's for running them from the unification scale to the scale relevant at colliders, and the bounds on the phases can get rather strong.

Nevertheless, since mass-spectra and couplings obviously depend on - besides several

¹bgaissma@ph.tum.de

real parameters - the phases, they are a priori non-negligible in studies of possible signals at future colliders and should be considered as free parameters of the model. Actually the situation is even worse: neglecting phases during the determination of real parameters from experimental data could lead to wrong inputs for theorists, who want to specify the underlying theory at the unification scale.

Despite the necessity to extract values for phases from experiment, the construction of sizable and comfortably accessible CP-violating observables is rather difficult in most of the production channels at an e^+e^- -machine as at least one secondary decay has to be included. Considering this problem we are introducing an object which quantifies the impact of non-vanishing phases on CP-even cross-sections and allows us to investigate this impact to some extent.

2 Idea, its Complementarity and Framework

The basic idea of this work [6] is to take today's low energy data (lower mass-bounds, d_e and a_μ) as a set of constraints for a parameter space scan and then to apply the resulting, low-energy compatible points to a set of high energy experiments @ NLC (e^+e^- and e^-e^- -option). This set of high energy-experiments is given by the total, unpolarized cross-sections for $\tilde{\chi}^0$ -, $\tilde{\chi}^-$ -, $\tilde{e}^-\tilde{e}^+$ pair production and is completed by $\tilde{e}^-\tilde{e}^-$ production [7],[8],[9],[10].

Since we are dominantly interested in the role of phases, the most stringent bounds on parameter space arise from the possible size of the SUSY-contributions to d_e and a_μ , which are given by:

$$\begin{aligned} 5.7 \times 10^{-11} < (a_\mu)_{\text{SUSY}} < 49.3 \times 10^{-11} \\ -13 \times 10^{-28} \text{ecm} < (d_e)_{\text{SUSY}} < 59 \times 10^{-28} \text{ecm} \end{aligned}$$

Recalling the diagrammatic structure of the SUSY-contribution to the leptonic dipole operators ($\tilde{\chi}^0\tilde{e}$ and $\tilde{\chi}^-\tilde{\nu}$) and comparing with the several (tree-level) diagrams for the given production modes we notice a *complementarity* between low- and high-energy observables: both low-energy observables always depend on products of different vertices while each diagram contributing to the cross-sections depends only on bilinears of one supersymmetric vertex (or a product of a supersymmetric one and a known SM-vertex). This implies that the low-energy observables connect different parts of SUSY-Lagrangians and therefore only can give bounds on combinations of phases, whereas high-energy observables can be used to investigate the different parts separately.

As framework for this project we are using the MSSM with R-parity, neglect sflavour mixing and assume horizontal universality of the Yukawa-like soft breaking A terms. Therefore the real parameters relevant for our analysis are:

$$|\mu|, \tilde{m}_L, \tilde{m}_R, |M_1|, M_2, |A|, \tan \beta,$$

and the investigated phases are ($\phi_2 = 0$ by convention):

$$\phi_\mu, \phi_1, \phi_A.$$

3 Significance and Mass Variation

As pointed out already, our aim is to find an object quantifying the impact of CP-odd phases on CP-even cross-sections and to understand the behaviour and meaning of this object. As underlying assumption for the definition of this object we assume that real parameters are already fixed. This assumption is indeed simplifying our analyses significantly, nevertheless we think it is a fair one, as most of the recent analysis do the same *and* neglect phases. The basic idea for the object we are introducing as *significance of a CP-even cross section with respect to CP-phases* is to compare the deviation of counting rates between a CP-conserving (CPC: real parameters, all phases $\equiv 0$) and CP-violating point (CPV: same real parameters, *but* phases $\neq 0$ and low-energy compatible) to the statistical error of the CPC-point. In formulae this idea reads as:

$$S \propto \frac{\Delta N_{CPV-CPC}}{\delta N_{CPC}}.$$

As there are two CP-conserving values ($0, \pi$) for each phase, we have to deal with 8 CPC points for each set of real parameters, a priori the same number of (different) significances is available for each cross-section and we have to decide which one to use in further discussions.

Our selection rule is then take the minimum of S with respect to these 8 points, this corresponds to the most conservative estimate of the impact of phases on cross-sections, and S finally reads:

$$S(\sigma_{f_i f_j}) = \min \left(\frac{|\sigma_{f_i f_j}^{CPV} - \sigma_{f_i f_j}^{CPC}|}{\sqrt{\sigma_{f_i f_j}^{CPC}}} \right) \sqrt{\mathcal{L}},$$

where the indices f_i, f_j refer a given out-mode with particles f_i, f_j and $\sigma_{f_i f_j}^{CPV/CPC}$ are the corresponding cross-sections in a CPV- or CPC-point.

Although it is rather clear that a high value of $S(\sigma_{f_i f_j})$ indicates a significant dependence of the mode on the phases, it is not clear where this dependence originates from. So the next step is to separate kinematical from coupling effects. For this purpose we define a variation of masses as:

$$V(m_i, m_j) = \frac{(m_i + m_j)^{CPV} - (m_i + m_j)^{CPC}}{(m_i + m_j)^{CPC}},$$

which is calculated in the CPV-point which minimizes $S(\sigma_{f_i f_j})$. Since slepton masses are phase independent we are investigating slepton production with respect to the lightest Neutralino in the t -channel diagrams. The point about studying this mass-variation is that a small variation of masses corresponds to small variation of the kinematical functions $(\beta(\frac{m^2}{s}), \lambda(\frac{m^2}{s}, \frac{m^2}{s}))$ and a large $S(\sigma_{f_i f_j})$ should therefore be induced by coupling effects.

4 Numerical Results

Our complete numerical analysis is currently performed for different values of $\tan \beta$ (3,6,9,12) and two values for $|\mu|$ (200 GeV, 500 GeV). The remaining real parameters are fixed as: $M_2=200$ GeV, $M_1=100$ GeV, $\tilde{m}_L=235$ GeV, $\tilde{m}_R=180$ GeV and $|A|=500$ GeV [6]. For each of these 8 real parameter points we scan the phases Φ_μ, Φ_1, Φ_A randomly in $(0, 2\pi)$, after applying the low-energy constraints as cuts the typical survival rate is a few permille (initially 500,000 points).

By correlating low-energy compatible values of the phases, we always find rather small bands for Φ_μ ($\mathcal{O}(\pi/6)$) located around 0, while the bands for Φ_1 and Φ_A can be rather large or even complete. The concrete pattern of correlations between them of course depends sensitively on the choice of real parameters. Anyway, these results are no news, so we don't present any of the low-energy correlations here.[4]

Although the number of correlations between low- and high-observables or between high-energy-observables is large, we can illustrate the basic results using two examples and a set of real parameters. As examples we show the correlations between d_e and the significance and the correlation between mass-variation and significance for the $\tilde{\chi}_1^0 \tilde{\chi}_2^0$ -mode (left and right, upper panels in Fig.1) and the $\tilde{e}_L^- \tilde{e}_R^+$ -mode (left and right, lower panels in Fig.1) in $e^- e^+$ -collisions.

Both modes illustrate that sizable significances are possible over the complete range of d_e , indicating possible big deviations between CPC and CPV-points even for vanishing d_e . The obvious no-correlation pattern between significances and d_e is rather transparent as the significances are functions of 2 phases while d_e depends on 3 phases. Note that d_e violates chirality and therefore always includes a factor of the electron Yukawa coupling, so that contributions proportional to A_e are *not* suppressed here. Basically the same argumentation applied to the strong correlation patterns between mass-variations and significances, as masses and cross-sections both depend on the same phases. Since the scale of mass-variation is of O(%) for both modes the significance must originate dominantly from coupling effects.

In Fig.2 we show the correlation between the significances of both modes, again we




Figure 1: upper boxes: $S(\tilde{\chi}_1^0 \tilde{\chi}_2^0)$ plotted against d_e (left panel) and $V(m_i, m_j)$ (right panel) for $|\mu|=200$ GeV and $\tan \beta=12$, lower boxes: same for $S(\tilde{e}_L^- \tilde{e}_R^+)$; $\sqrt{s}=500$ GeV and $\mathcal{L}=500 \text{ fb}^{-1}$.

observe a correlation pattern as both cross-sections depend on the same phases. The triangle structure of the correlation is clear, as the significance of each mode reaches a maximum for a certain value of Φ_1 (Φ_μ is a small band and can be considered as fixed) and the maximizing values for different cross-sections do *not* coincide.




Figure 2: Correlation between $S(\tilde{e}_L^- \tilde{e}_R^+)$ and $S(\tilde{\chi}_1^0 \tilde{\chi}_2^0)$ for $|\mu|=200\text{GeV}$ and $\tan\beta=12$ for $\sqrt{s}=500\text{ GeV}$ and $\mathcal{L}=500\text{ fb}^{-1}$.

5 Conclusion

After using low-energy data as constraints on parameter space we rediscover older bounds on phases, the allowed range for the phases can be rather big. We can give a quantity (*significance*) to estimate the impact of CP-odd phases on CP-even cross-sections. This significance can be rather big, implying a significant deviation of CP-violating scenarios from CP-conserving one already at the level of cross-sections. These significances are correlated with each other, *but not* with d_e and a_μ and they are dominantly due to coupling effects. Therefore phases should be taken into account in any discussion of how to extract model parameters from measured cross-sections.

Acknowledgements

The project is the result of a collaboration with S.Y.Choi and M.Drees. The work of BG is supported in part by the Deutsche Forschungsgemeinschaft, project number DR 263.

References

- [1] D. E. Groom *et al.* [Particle Data Group Collaboration], Eur. Phys. J. C **15**, 1 (2000).
- [2] G. W. Bennett *et al.* [Muon g-2 Collaboration], Phys. Rev. Lett. **89**, 101804 (2002) [Erratum-ibid. **89**, 129903 (2002)] [arXiv:hep-ex/0208001].
- [3] M. Hayakawa and T. Kinoshita, arXiv:hep-ph/0112102.
- [4] M. Brhlik, G. J. Good and G. L. Kane, Phys. Rev. D **59**, 115004 (1999) [arXiv:hep-ph/9810457].
- [5] T. Ibrahim and P. Nath, Phys. Rev. D **58**, 111301 (1998) [Erratum-ibid. D **60**, 099902 (1999)] [arXiv:hep-ph/9807501].
- [6] S. Y. Choi, M. Drees and B. Gaissmaier, in preparation
- [7] S. Y. Choi, J. Kalinowski, G. Moortgat-Pick and P. M. Zerwas, Eur. Phys. J. C **22**, 563 (2001) [Addendum-ibid. C **23**, 769 (2002)] [arXiv:hep-ph/0108117].
- [8] S. Y. Choi, A. Djouadi, M. Guchait, J. Kalinowski, H. S. Song and P. M. Zerwas, Eur. Phys. J. C **14**, 535 (2000) [arXiv:hep-ph/0002033].
- [9] M. E. Peskin, Int. J. Mod. Phys. A **13**, 2299 (1998) [arXiv:hep-ph/9803279].
- [10] S. Thomas, Int. J. Mod. Phys. A **13**, 2307 (1998) [arXiv:hep-ph/9803420].